

Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

Sper I. ps. 864 Choisy monograp, 179 Nº 16 Cuscula chilensis Her Dasella mbm (hvis)? ex Herb In A. Jong New York Coule ramoso, floribus subsessilibus glomentis ompalities i tubo corollac cylindrico calycis cupuliti campanulati segmenta rotundata obtusa murulatas et lacricias limbi ovalas obtusas palentes (demun reflexas) triple superante, of antheris sessility inclusis; squamis ovitis margine levers, Carries iteraries subinouvis; stylis ovanium conicum stylopodio coronatura superantitus, stigmations oupitalis, inclusis - capsuls Flowers have at lange as the largest Morth ancer's Constante only campanulate, not adpressed to the Corolla shis is the case in all the other Cosmitas It know for the Cales handly reached up to the middle of the Cong orghindre thele This South america plant also belongs to The Unite Anen rance stigmatibus capitation! Sprangel C. C. says; corolla urreolata, best my spensmen has a corolla cylindria. Though MISSOURI

BOTANICAL copyright reserved cm GARDEN


Clematis Vitirella La favorata Pers for IT, 865 AII 308 CEORGE ENGELMANN PAPERS MISSOURI BOTANICAL GARDEN A.B. Fo 1820 Europe med et aust. Af mohr. MISSOURI BOTANICAL copyright reserved cm GARDEN

Cuscuta corymbosa R. a Par. - Cho. 29! Caule subrapillares vamos, fascriralis floren subremilibus, compositis, floribus singulis pedstellati, 5 partitis, onlyres semigurguefile. bobis obhesius ratio, tato vorollee campa; mulato turbinito algreen duplo excedente Cerinis limbi odatis acutes aprice inflexis patentibus sub duplo longiore. stamphotus Cialo mulle brevior bus filamente a basi let, subulti antheris oblogs - Circan bus, squaming breviter frubrites, fruiten conolla pyriforme agreentibus francen corolla attrigentibus, stigmatibus, organitations On Medingo-sative Tonilis ets. Geneva Inch for South amen on Choisy This plant is the same as the Chamiara, though the flowers are (at least in my sperimens) hat auch more than half as large, there shorten are some Ther distinctions, experially the rowlle is a little the lobes of the conthe are langly Bilital at base and imbritated the rules are larger i proportion and not depty laverated but rathe shortly frubricted in and the whole texture of the parts of the flower is firmer even somewhat gland land land the printer of human tenstity; shape of take of corolla (anthurs), flaments) inflaments) 2 it Aug & Savon's Runt ? In one flower the scales were nearly wanting only along some of the short flament, shere they athere to the corolle namour or widow wings oned be observed which explain the notice of these rockes as well as the normal MISSOURI

BOTANICAL

GARDEN

copyright reserved



Coscula hassiaia Meist Carle filiporas varnoso, fasoriralis florum subsessilibres. lexis tabapositis, floribus singulis · longues pedirelletis, 5 partites, * tubo alyses semiguinquefide, lobis corollae turbinato (obronico) calgrem mullo excedente, lagrionies limbs ountes arumine inflexo cuspidatos sub duplo * erestis s. petentitos longione, staminibres limbo multe brevionibres filamentes e lassi let, subulation anthen, oblonge - linear bus ; squamis finbriata leveris, faurem the corollar non attinguntions (lasi carnosse constructo suffellum) acquestitus fancem corolle attengentibres. strymetetus opertalis. Capsula in Near Casel, Hassin and in other party of thirte Germany on Medingo sation Meleteter Anthemin Podospermen Bakhansin Somehus, Galican St., Alexan Graminese n Medryo (most Rudio). ____ Impater for Bett america? This interesting species has been for the fell 1843 for the first time as at different places is with a Garmany by or ferrent observers. The spermen desontentes en has been collected near Counce of In Mitte Tu flowers are and the very oformer. IL van hundly been doubled, that (Austi this glessies with reportate stymes has not been intendered for Both to his long South amenda. It hear a great The my ments of the only are not interinted, non It is remarkable by the teerbinate is the base of the lacinine of the rowler white founded founded founded founded founded founded at here and widens towards the By the select or hooks thought by the slongated somewhat points of the other se linear anther and by the ovary obtuse jutals which has a fleshy rate men their mer love and appears therefore pear shaped This love of the ovary appears to be just the reverse of the thurthere of many north amening openies (especially C. gloments) so MISSOURI

BOTANICAL copyright reserved GARDEN

a flesky stylopodium is suppeningosed on the capsule. The flowers are under the texture of a thin texture of larger the any x atte of laments are conspien are most north amens easily recognized along species they are probable near there shale commercial with C. Chilensis is size Moveragh the the lake of the corolla shape be different. Thapsale I have not seen but suppose for The appearance of a few flowers gird after flowers, that the tuke will som become ventre voze, and the apple of on amount to of the sipening expende It is singular that handy any no ripe seeds evene obteried Jennany which wild the ment is a fact favourable to the supp of its being introduced MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS MISSOURI BOTANICAL


copyright reserved cm GARDEN


Lepidanche putrisur adpresso Repridanshe futicum aule ramoso elato posibis sessilibre vapralis 1842
glomentis o parti tis the roulle learne cremulations ogladio calquem 7 guarum policies adpressio total interior production ex reducte mate about the stand Carinias limbs oblonges, obtusas mitter exterior bus putentes leis superante; staminibus ovato ovorbulation verse ovater, ikkenbrit, link brevstribus, squamis mintiple lavinitis convergentions ovann in ludentibus. ovario om stylopeli stylo, requente, capsula globon subscrite marrescente obtesta, 2 = 4 Juna Asserved this species in System of the said of the Manus amongst a most leek an ant groth of river and modern but of Afferent that grown over Organie, Mus, their 2 Wit. Comus Lum Stengon and others, all of them the stand plants that the western Virginia AG. & VES. Sept. 1843 The the land Cennlas nearer than the the Theries of this Jenus in habit as well as Im the temeller muchen of sophs the adjuncted algorine trales The flowers are dosely serile but distinct a with as in a composition where it is hardly possible to Istryunsh a count

Asker pennicenes L hur. A. amoenus Lam hispidus Lam Jac.11/533 Firmus De MISSOURI BOTANICAL GARDEM GEUNGE ENGELMANN PAPERS Amer. Bor. H. Jalyw 1825 MISSOURI BOTANICAL copyright reserved cm

GARDEN

Sportante fatilie the single glowers. The glomentes cither form a continues live rough Sten of the parent plant one they are separate more como this flips are white to form strong patty 18.2) stems are white to for bally for the shade i which the plant grows at the flowers are at the flowering time, as those of a compositan. The raps les are generally 2 suder but as they arend no enoused in it the the pures they are also from 3 to 4 seeded





Lepidanche compacta Just Lepidanche compacta ">p. Legisdanske tomprette Caule ramoso, elato, floribus sessilibus gomentis, 5- partitis, sepulis sub novem leviter armelites convavis adpressis orbinlates interioribus minoribus, tube molle cylindrio, colycon colycon Int septemento, orlynem red lamins limbi Cinani oblinger obtersay Duplo superante. James, Lam Links brevion bus . squaring. primalified lacinistis, cranio oum styloperd styler saguit. capsula globon subante corolle marressente oblevta 1-4 sperma Mailie of Sitt Caroline S. D. Darkley, 1842 Monthis of the state worth Carolin Saller South 1843 Sides of Negro mustain N.C. and on Almus Confly wither - Alebania, om Sornor glaber, Herb. An Grug. This is priced the Customers of Offabre & George The State of the State of George of the State o represent a peril the L. adjuster which they for has only been for on the bottom to 3 gluen tanks will of the Mississiff and allihoris noen It is very remakette that again in this instance two very nearly selle speries from the one for the South Allegany Munters and the the western Low land, analogie, offen Baptisin alla & Befin aumanth Pharelin finbrit & Much Curshi! a Soller. Star Months from The whole appearing of the solution of the western valutive its of former of the state its of former of the state of t MISSOURI


BOTANICAL copyright reserved cm GARDEN


TurI 113 Lavandula abrofanoides dans A. PHALET Ins. Enar. MISSOURI BOTANICAL GARDERS MISSOURI BOTANICAL

copyright reserved

GARDEN

Lysbanie computette of the corolla excreed the close a compact scales of the caly considered and is much theman in properties Eits length. It gives then for to the capsule, which it covers! Ethough the composale itself is nearly globose f. At is the to most early Stringer to form of the points fruit parmit om spiries to be more easily and immediately dishinguishe for a morrow, just after Monening. The corolla appears to be more membranercous than in the wester species and remark, estelish, she dry, the other would turno redsish Born.





Cusetta glomente Choisy L'épidanthe Compositann L'Emporitanun caule ramoso humi li', floritus

sassi, libra aurite ronglomentis quamis ralyrinis 10-15 inhisti;

5 partitis tule rorolla drenn latis onspir apire

solveliadris ralyren succes reflexis, interioristis majoris aquintation excedente leninas l'ubi obloques oblans matentes o. reflexas les sepande, Marianis Combum agreentiber, extally James pohoulofich lariate convergentibus, ovanium obte includation oranio skylopodrio skyles 2 Freviore, rupsula globosa, squami relymini shvoler nat. Corolle marres rente colecté! 1-2 // ma telylopodio oun stylis consult d. L. Solden fortus minoring levinis torble reflex is stylopodie ovans de eminore B. Lelianthi florten mejorbes, lankis voolle patention, sotalis, squaris
inter se connatis coronan formantion, style jodie ovario majore This singular plant appears to be peculiar to the wintern I have observed it this styden is I mairied plant since (833 in west soil prairies vone Modernie Grier on Silphium & Clapp = Juliana Co Clapp = Juliana genety on Helather since 1858 - pinniles Corne; Ties, flowing - Any & Lent. A seneme Description of Their habit MISSOURI

BOTANICAL

GARDEN

copyright reserved

Dens. I, 316 Roch (II, 211 Spr I, 905 Lium repens L. Un pratis humidis Vinundatis German. austr. Anglia Vania. In paludosis nemorosis prope Darmsladt Grieshem, Gerau 1826

0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

GARDEN

MISSOURI
BOTANICAL
GARDEN

L'apportann 2 harben given above when nuky of the genus. Soth vanisties may prove Distinct Species but for the present I am inable to Distryuis the by more , hopeortant characters The flower on always spected the take is not exactly Eylander The flowers on always spected the take wider at the mouth the Myles one longer than in the late the lase, rather obcorries. any Men of our discretas, and gener nearly always unique the Plyms is reprétate asi all the Amonin assentes, the somste and are viseled on Anding Shylopodium asia which is layer than in any and this is more remarks - the Locivaginis that i lelianthi, when the shape is nearen the oglind, sol on. stems ved, filipon, brushoj, flows white (dug. 1843) This Speries is before the buying of August entirely similar to any othe Cusante, and covers something with its numerous red or orange slams in bounder, whole petities in low processes, so that they are be seen few off. About that time The stems where they closely embrace the stem of the moth plant would become flettened, swell on the sides in irregular lines or knots; and soon the expressions bursts there and now the little rounded squarrose leafy flowerbuts make their appearen and grow rapidly so that the Tur Cateral Chy som disappen at The whole are manged in the thork and orowded ropelike glomentes are alongs Ishickal. - Joursome time the Caps are covered at They been by the bown rays of the existences of thisken. MISSOURI

BOTANICAL

GARDEN

copyright reserved

